

Gill-Thwaites, H., Elliot, K., & Morrissey, A-M. (2021). LOCCATE: A tool to identify the diagnostic spectrum profile of motor function and functional communication responses for the individual with a prolonged disorder of consciousness. *Neuropsychological Rehabilitation*. doi: 10.1080/09602011.2021.1981949

Helen L. Gill-Thwaites, Karen E. Elliott & Anne-Marie Morrissey (2021): LOCCATE: A tool to identify the diagnostic spectrum profile of motor function and functional communication responses for the individual with a prolonged disorder of consciousness, *Neuropsychological Rehabilitation*, DOI: 10.1080/09602011.2021.1981949

H. Gill-Thwaites, K. E. Elliott & R. Munday (2017): SMART – Recognising the value of existing practice and introducing recent developments: leaving no stone unturned in the assessment and treatment of the PDOC patient, *Neuropsychological Rehabilitation*. DOI 10.1080/09602011.2017.1310113

Tennant A. and Gill-Thwaites H. (2016). A study of the internal construct and predictive validity of the SMART assessment for emergence from vegetative state. *Brain Injury*, 1–8. Published online. DOI: 10.1080/02699052.2016.1225983.

da Conceição Teixeira L., Gill-Thwaites H., Reynolds F., et al. (2016). Can behavioural observations made during the SMART assessment detect the potential for later emergence from vegetative state? *Neuropsychological Rehabilitation*, 1–10. DOI:10.1080/09602011.2016.1243482.

Gill-Thwaites H. and Munday R. (2010). The Sensory Modality Assessment and Rehabilitation Technique (SMART): a comprehensive and integrated assessment and treatment protocol for the vegetative state and minimally responsive patient. *Neuropsychological Rehabilitation*, 9: 305–320.

Gill-Thwaites H. (2006). Lotteries, loopholes and luck: misdiagnosis in the vegetative state patient. *Brain Injury*, 20: 1321–1328.

Gill-Thwaites H. and Munday R. (2004). The Sensory Modality Assessment and Rehabilitation Technique (SMART): a valid and reliable assessment for the vegetative and minimally conscious state patients. *Brain Injury*, 18: 1255–1269.

Wilson S.L. and Gill-Thwaites H. (2000). Early indication of emergence from vegetative state derived from assessment with the SMART: a preliminary report. *Brain Injury*, 14: 319–331.

Gill-Thwaites H. (1997). The Sensory Modality Assessment Rehabilitation Technique: a tool for assessment and treatment of patients with severe brain injury in a vegetative state. *Brain Injury*, 10: 723–734.

Wilson S.L., Brock D., Powell G.E., et al. (1996). Constructing arousal profiles for vegetative state patients: a preliminary report. *Brain Injury*, 10: 105–113.

Wilson S.L., Powell G.E., Brock D., et al. (1996). Vegetative state and responses to sensory stimulation: an analysis of 24 cases. *Brain Injury*, 10: 807–818.

Wilson S.L., Powell G.E., Elliott K., et al. (1991). Sensory stimulation in prolonged coma: four single case studies. *Brain Injury*, 5: 393–400